

Several Limit Theorems on Fuzzy Quantum Space

Viliam Ďuriš^{1,*}, Renáta Bartková² and Anna Tirpáková^{1,3}

¹ Department of Mathematics, Constantine The Philosopher University in Nitra, Tr. A. Hlinku 1, 949 74 Nitra, Slovakia; atirpakova@gmail.com

² Podravka International s.r.o, Janka Jesenského 1486, SK-960 01 Zvolen, Slovakia; renata.hanesova@gmail.com

³ Department of School Education, Faculty of Humanities, Tomas Bata University in Zlín, Štefánikova 5670, 76000 Zlín, Czech Republic

* Correspondence: vduris@ukf.sk; Tel.: +421-37-6408-708

Abstract: The probability theory using fuzzy random variables has applications in several scientific disciplines. These are mainly technical in scope, such as in the automotive industry and in consumer electronics, for example, in washing machines, televisions, and microwaves. The theory is gradually entering the domain of finance where people work with incomplete data. We often find that events in the financial markets cannot be described precisely, and this is where we can use fuzzy random variables. By proving the validity of the theorem on extreme values of fuzzy quantum space in our article, we see possible applications for estimating financial risks with incomplete data.

Keywords: fuzzy quantum space; convergences on fuzzy quantum space; law of large numbers; central limit theorem; Fisher–Tippett–Gnedenko theorem; Balkema; de Haan–Pickands theorem



Citation: Ďuriš, V.; Bartková, R.; Tirpáková, A. Several Limit Theorems on Fuzzy Quantum Space. *Mathematics* **2021**, *9*, 438. <https://doi.org/10.3390/math9040438>

Academic Editor:
Gustavo Santos-García

Received: 2 February 2021
Accepted: 18 February 2021
Published: 23 February 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Selected limit theorems, which we shall deal with in the article, are well known from Kolmogorov's classical probability theory. Kolmogorov's work [1] introduced the theoretical axiomatic model in which events connected with the experiment form the Boolean σ -algebra of subsets \mathcal{S} of the set Ω . Thus, the probability is that for the σ -additive, nonnegative final function P on \mathcal{S} , with values in the interval $[0, 1]$, if $\{A_n\}$ is a sequence of mutually exclusive events from \mathcal{S} , then $P(\cup_n A_n) = \sum_n P(A_n)$ and $P(\Omega) = 1$. Limit theorems have a wide range of use in this theory. Their validity has already been proven for other structures (spaces), e.g., MV-algebras defined in [2]. We want to extend their use; therefore, in this article we prove that they also apply to sets in which we are working with incomplete data. Specifically, they also apply to Fuzzy quantum space, and that is the most significant finding in this article.

After some time, it became apparent that Kolmogorov's classical model of the probability theory was not sufficient for describing quantum mechanics situations. Birkhoff and von Neumann [3] referred to the fact that the set of experimentally verifiable statements about the quantum mechanical system does not have the same algebraic structure as Boolean algebra. Heisenberg [4] and Schrödinger [5] put forth the earliest attempts at the mathematical formulation of quantum mechanics. Schrödinger presented the formalism of wave mechanics, while Heisenberg proposed the formalism of matrix mechanics.

Zadeh [6] wrote about the theory of fuzzy sets in the 1960s. The current quantum theory, basic mathematical model is that of von Neumann, grounded in the geometry of Hilbert space (Varadarajan, [7]). If we define all closed subspaces of a given Hilbert space (where, according to Varadarajan, the notion "a state of system" means a measure of probability on \mathcal{M}) as system \mathcal{M} , and such a definition is compared with that of the P -measure on fuzzy sets (according to Piasecki [8]), it follows that both objects have a similar algebraic structure. Piasecki submitted a model called soft σ -algebra in the fuzzy set theory in 1985. His model demonstrated several characteristics identical to quantum

logics. That comparison was first noted by Riečan [9] and then by Pykacz [10], and it led us to the idea to build a quantum theory based on fuzzy sets. If A is a non-empty set called a universum and \mathcal{M} is a system of fuzzy subsets of universum A , i.e., the system of functions on A with values in the interval $[0, 1]$, then according to Riečan [9] we say that (A, \mathcal{M}) is an F -quantum space, also referred to by Dvurečenskij and Chovanec [11] as a fuzzy quantum space, or by Dvurečenskij [12] as a fuzzy measurable space.

Many writers have attempted to prove some known assertions from the classical probability theory in the theory of fuzzy quantum spaces. For example, Dvurečenskij [12], Navara [13], and Navara and Pták [14,15] studied the existence of a fuzzy state on fuzzy quantum space while Dvurečenskij and Riečan [16,17] examined joint fuzzy observables and joint distributions of fuzzy observables. The representation theorem was proved by Dvurečenskij, Kôpka, and Riečan [18]; it also includes the case in fuzzy quantum space. Riečan [19,20] looked at the theory of an indefinite integral on fuzzy quantum space. Mesiar [21–23], Piasecki [24,25], and Piasecki and Svitalski [26] investigated the extension of the validity of the Bayes formula for fuzzy sets. Markechová [27–29] researched the entropy on fuzzy quantum space, and Tirpáková and Markechová [30] investigated the fuzzy analogies of some ergodic theorems and Birkhoff's individual ergodic theorem and maximal ergodic theorem for fuzzy dynamical systems [31].

The existence of the sum of fuzzy observables is a key fact for the analysis of many assertions in the fuzzy sets theory. The existence of the sum of compatible fuzzy observables was proved by Harman and Riečan [32].

Among the vital concepts of probability theory are the different kinds of convergence of random variables. They are especially significant for parts dealing with the validity of various forms of the law, the central limit theorem, and big numbers. As a result, the problem of generalizing different types of convergence for fuzzy quantum space (A, \mathcal{M}) became topical. A few authors studied particular types of convergences on quantum logic. Here, we mention the writings which were the basic material for the study of various types of convergences of fuzzy observables on fuzzy quantum space (A, \mathcal{M}) : Dvurečenskij and Pulmanová [33], Jajte [34], Ochs [35,36], Cushen [37], Gudder [38], and Révész [39]. Some types of convergences of fuzzy observables on fuzzy quantum space were dealt with by Dvurečenskij [40], Riečan [41,42], Chovanec and Kôpka [43], Kôpka and Chovanec [44], and others.

We formulated convergences and consequently proved many familiar limit theorems [34,36,45] for fuzzy quantum space on the basis of the analogy of the probability theory notions. As the central limit theorem refers to the limit distribution of the averages of independent, equally-distributed random variables, extreme value theory (EVT) addresses the limit distribution of the maximums of the independent, equally-distributed random variables [45,46]. EVT's principal objective is to know or predict the statistical probabilities of events that have never or rarely been observed. Kotz and Nadarajah [47] indicated that the extreme value distributions could be traced back to Bernoulli's 1709 work [48]. The theory of max-stable distribution functions, the counterpart of Feller stable distributions [49], formed the basis of the probability background. First, the statistical analysis of extreme values was performed in order to study flood levels. These days, the areas of application include finance, meteorological events, insurance, industry, or the environmental sciences [50]. Allow X_1, X_2, \dots, X_n to be a sequence of n , independently and identically distributed random variables with distribution function F . The corresponding ordered sequence in non-decreasing order is indicated by $X_{1:n}, X_{2:n}, \dots, X_{n:n}$, where $X_{i:n}$, $i = 1, \dots, n$ represents the i -th order statistic. $X_{1:n}$ and $X_{n:n}$ stand for the sample minimum and the sample maximum, respectively. Then, examine the sequence of maxima $M_1 = X_1$, $M_n = X_{n:n} = \max(X_1, X_2, \dots, X_n)$, for $n \geq 2$, obtained from the above sequence. All the sequence minimum results can be obtained from those of the sequence maximum since $m_n = \min(X_1, X_2, \dots, X_n) = -\max(-X_1, -X_2, \dots, -X_n)$. M_n 's exact distribution can be obtained from the distribution function F . In fact, for all $x \in \mathbb{R}$:
$$F_{M_n}(x) = P(M_n \leq x) = P(X_1 \leq x, X_2 \leq x, \dots, X_n \leq x) = \prod_{i=1}^n P(X_i \leq x) = F^n(x).$$
 For a

single process, the behavior of the maxima can be described by the three extreme value distributions: Gumbel, Fréchet and reversed Weibull distribution as suggested by the Fisher–Tippett–Gnedenko theorem. One can combine these three distributions into a single family of continuous cumulative distribution functions, known as the generalized extreme value (GEV) distributions [50]. A GEV can be identified by the real parameter γ and the extreme value index, and as a stable distribution it is a characteristic exponent $\alpha \in [0, 2]$. Subsequently, several researchers have provided useful applications of extreme value distributions. They may be found in several works [51–54].

2. Fuzzy Quantum Space

First, we recall the definitions of basic notions and some facts that will be used in the following text. In the quantum space approach to the fuzzy quantum theory, the triple (Ω, S, P) is replaced by the couple $(\mathbb{A}, \mathcal{M})$ where \mathbb{A} is a nonempty set, $\mathcal{M} \subset [0, 1]^{\mathbb{A}}$ is fuzzy σ -algebra of fuzzy subsets of \mathbb{A} , such that the following conditions are satisfied:

- (i) if $1_{\mathbb{A}}(x) = 1$ for any $x \in \mathbb{A}$, then $1_{\mathbb{A}} \in \mathcal{M}$
- (ii) if $f \in \mathcal{M}$, then $f'1_{\mathbb{A}} - f \in \mathcal{M}$
- (iii) $\bigvee_{n=1}^{\infty} f_n \sup_n f_n \in \mathcal{M}$, for any $\{f_n\}_{n=1}^{\infty} \subset \mathcal{M}$
- (iv) if $\frac{1}{2_{\mathbb{A}}}(x) = \frac{1}{2}$ for any $x \in \mathbb{A}$, then $\frac{1}{2_{\mathbb{A}}} \notin \mathcal{M}$

Elements of the set \mathcal{M} are called fuzzy subsets of the universe \mathbb{A} . In particular, if f is the characteristic function, we call it a crisp set. The symbols $\bigvee_{n=1}^{\infty} f_n \sup_n f_n$ and $\bigwedge_{n=1}^{\infty} f_n \inf_n f_n$ indicate a fuzzy union and a fuzzy intersection of the sequence of fuzzy sets $\{f_n\}_{n=1}^{\infty} \subset \mathcal{M}$, respectively. The event $f'1_{\mathbb{A}} - f \in \mathcal{M}$ is the so-called fuzzy complement. By Piasecki [8], the system \mathcal{M} is called a soft σ -algebra.

To define and prove the law of large numbers and the central limit theorem, we need the following basic notions:

Definition 1. A fuzzy state on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ is a mapping $m : \mathcal{M} \rightarrow [0, 1]$, such that

- (i) $m(f \vee (1_{\mathbb{A}} - f)) = 1$ for every $f \in \mathcal{M}$
- (ii) if $\{f_k\}_{k=1}^{\infty}$ is a sequence of pairwise orthogonal fuzzy subsets from \mathcal{M} , i.e., $f_i \perp f_j$, $(f_i \leq 1_{\mathbb{A}} - f_j)$, whenever $i \neq j$, then $m(\bigvee_{k=1}^{\infty} f_k) = \sum_{k=1}^{\infty} m(f_k)$

According to Piasecki [8], a fuzzy state is called the σ -measure. The triplet $(\mathbb{A}, \mathcal{M}, m)$ where m is a σ -measure is called a fuzzy probability space. This structure was studied in [55,56].

For illustration, we give the following example of a nontrivial fuzzy quantum space [55].

Example 1. Consider $(\mathbb{A}, \mathcal{M})$ where $\mathbb{A} = [0, 1]$, $f : \mathbb{A} \rightarrow \mathbb{A}$, $f(x) = x$, $\mathcal{M} = \{f, f', f \vee f', f \wedge f', 0_{\mathbb{A}}, 1_{\mathbb{A}}\}$ for every $x \in \mathbb{A}$. It is evident that $f \vee f' \neq 1_{\mathbb{A}}$. We define the mapping $m : \mathcal{M} \rightarrow [0, 1]$ by the equalities $m(1_{\mathbb{A}}) = m(f \vee f') = 1$, $m(0_{\mathbb{A}}) = m(f \wedge f') = 0$, and $m(f) = m(f') = \frac{1}{2}$. Then the triplet $(\mathbb{A}, \mathcal{M}, m)$ is a fuzzy probability space.

Definition 2. A fuzzy observable on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ maps to $x : \mathcal{B}(\mathbb{R}^1) \rightarrow \mathcal{M}$, satisfying the following properties:

- (i) $x(E^c) = 1_{\mathbb{A}}(x) - x(E)$ for every $E \in \mathcal{B}(\mathbb{R}^1)$
- (ii) if $\{E_n\}_{n=1}^{\infty} \subset \mathcal{B}(\mathbb{R}^1)$, then $x(\bigcup_{n=1}^{\infty} E_n) = \bigvee_{n=1}^{\infty} x(E_n)$

where $\mathcal{B}(\mathbb{R}^1)$ denotes the Borel σ -algebra of the real line \mathbb{R}^1 and E^c denotes the complement of a set E in \mathbb{R}^1 .

Definition 3. Let $f \in \mathcal{M}$. The mapping $x_f : \mathcal{B}(\mathbb{R}^1) \rightarrow \mathcal{M}$ is defined by

$$x_f(E) = \begin{cases} f \wedge f', & \text{if } 0, 1 \notin E \\ f', & \text{if } 0 \in E, 1 \notin E \\ f, & \text{if } 0 \notin E, 1 \in E \\ f \vee f', & \text{if } 0, 1 \in E \end{cases}$$

for every $E \in \mathcal{B}(\mathbb{R}^1)$ there is a fuzzy observable of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ called the indicator of fuzzy set $f \in \mathcal{M}$.

In particular, the null fuzzy observable of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ maps to $o : \mathcal{B}(\mathbb{R}^1) \rightarrow \mathcal{M}$ defined by

$$o(E) = \begin{cases} 0_{\mathbb{A}}, & \text{if } 0 \notin E \\ 1_{\mathbb{A}}, & \text{if } 0 \in E \end{cases}$$

where $E \in \mathcal{B}(\mathbb{R}^1)$.

If $\tau : \mathbb{R}^1 \rightarrow \mathbb{R}^1$ is a Borel measurable function and x is a fuzzy observable, then $\tau \circ x : E \rightarrow x(\tau^{-1}(E)), E \in \mathcal{B}(\mathbb{R}^1)$ is a fuzzy observable, too. In this way, we define the functional calculus of fuzzy observables. For example, if $\tau(t) = t^2, t \in \mathbb{R}^1$, we write $\tau \circ x = x^2$ and the like. In particular, if $a \in \mathbb{R}^1$, then $ax : E \rightarrow x(\{t \in \mathbb{R}^1 : at \in E\})$ for any $E \in \mathcal{B}(\mathbb{R}^1)$.

Let x be a fuzzy observable of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ and let $B_x(t) = x((-\infty, t)), t \in \mathbb{R}^1$. Dvurečenskij and Tírpáková [57] proved that the system $\{B_x(t) : t \in \mathbb{R}^1\}$ of fuzzy sets of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ is a one-to-one correspondence to fuzzy observable x . Due to this result, the sum of any pair x and y of fuzzy observables of $(\mathbb{A}, \mathcal{M})$ can be introduced as follows:

Definition 4. Let x and y be two fuzzy observables of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$. If the system $\{B_{x+y}(t) : t \in \mathbb{R}^1\}, B_{x+y}(t) = \bigvee_{r \in \mathbb{Q}} (B_x(r) \wedge B_y(t - r)), t \in \mathbb{R}^1$, where \mathbb{Q} is the set of all rational numbers, then we determine fuzzy observable z of $(\mathbb{A}, \mathcal{M})$. We call it the sum of x and y and write $z = x + y$.

In the text [57], it was proved that the sum of two observables always exists, and it coincides with the pointwisely-defined sum of observables for σ -algebra of crisp subsets. Moreover, $x + y = y + x, (x + y) + z = x + (y + z)$ for fuzzy observables x, y , and z . The subtraction of fuzzy observables x and y is defined as follows: $x - y = x + (-y)$, where $(-y)(E) = y(\{t : -t \in E\}), E \in \mathcal{B}(\mathbb{R}^1)$. The mean value of a fuzzy observable on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ was defined by Riečan [58] as follows: Let x be a fuzzy observable, and let m be a fuzzy state. If the integral $m(x) \int_{\mathbb{R}^1} t dm_x(t)$ exists, then $m(x)$ is called the mean value of x in m , where $m_x : E \rightarrow m(x(E)), E \in \mathcal{B}(\mathbb{R}^1)$ is a probability measure on $\mathcal{B}(\mathbb{R}^1)$. In addition, if u is a Borel measurable function, then $m(u \circ x) \int_{\mathbb{R}^1} u(t) dm_x(t)$ in the sense that if one side exists, then the second side exists too, and they are equal. Specially, if $u(t) = (t - m(x))^2$, then $D(x)m((x - m(x))^2)$ is called the dispersion of fuzzy observable x in fuzzy state m .

3. Convergences on a Fuzzy Space

Various types of convergences of random variables belong among important concepts of the probability theory. Therefore, the notion of a fuzzy observable is an analogy to the notion of a random variable. When defining different types of convergence and for the proof of limit theorems on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$, we used the method of F - σ -ideals, which enabled us to reformulate and prove many of the known limit theorems of the classical probability theory for the fuzzy quantum space $(\mathbb{A}, \mathcal{M})$. The basic idea of the F - σ -ideals method is described in [54], and we can shortly describe it as follows: Let m be a fuzzy state on fuzzy

quantum space $(\mathbb{A}, \mathcal{M})$. Denote $I_m = \{f \in \mathcal{M} : m(f) = 0\}$. Dvurečenskij and Riečan [16] proved that I_m is a σ -algebra of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$. The relation “ \sim_m ” defined on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ via $f \sim_m g$ if and only if $m(f \wedge g') = 0 = m(g \wedge f')$ is the congruence, and, moreover, $\mathcal{M}/\sim_m = \{\bar{f} = \{g \in \mathcal{M} : g \sim_m f\} : f \in \mathcal{M}\}$ is the Boolean σ -algebra (in the sense of Sikorski [59]), where complementation “ $'$ ” in \mathcal{M}/\sim_m is defined with properties $(\bar{f})' = \overline{f'}$, $f \in \mathcal{M}$ and $\bigvee_i \overline{f_i} \bigvee_i \overline{f'_i}$, $\bigwedge_i \overline{f_i} \bigwedge_i \overline{f'_i}$, $f_i \in \mathcal{M}$). Then, according to these properties, the mapping $h : \mathcal{M} \rightarrow \mathcal{M}/\sim_m$ defined by $h(f) = \bar{f}$, $f \in \mathcal{M}$ is a σ -homomorphism from \mathcal{M} onto \mathcal{M}/\sim_m . The mapping μ from a Boolean σ -algebra \mathcal{M}/\sim_m into the interval $[0, 1]$, defined by $\mu(\bar{f}) = m(f)$ for every $\bar{f} \in \mathcal{M}/\sim_m$, is a probability measure on the Boolean σ -algebra \mathcal{M}/\sim_m . According to the Loomis–Sikorski theorem in [60], there is a measurable space (Ω, \mathcal{S}) and σ -homomorphism φ from \mathcal{S} onto \mathcal{M}/\sim_m , and due to Varadarajan [7] there are functions $u, u_1, u_2, \dots : \Omega \rightarrow \mathbb{R}^1$, such that

$$\varphi(u_i^{-1}(E)) = h \circ x_i(E), i = 1, 2, \dots, E \in \mathcal{B}(\mathbb{R}^1) \tag{1}$$

$$\varphi(u^{-1}(E)) = h \circ x(E) \tag{2}$$

where $h \circ x$ is an observable of a Boolean σ -algebra \mathcal{M}/\sim_m . Moreover, mapping $\mu_\varphi : \mathcal{S} \rightarrow [0, 1]$, defined as $\mu_\varphi(\Lambda) = \mu(\varphi(\Lambda))$, $\Lambda \in \mathcal{S}$, is a probability measure on \mathcal{S} .

Gudder and Mullikin [51] introduced many types of convergences for observables in quantum logics. Inspired by their definition, Dvurečenskij and Tirkaková introduced [57] the following definition:

Definition 5. We say that sequence $\{x_n\}_{n=1}^\infty$ of fuzzy observables on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ converges to fuzzy observable x

(i) in fuzzy state m , if for every $\varepsilon > 0$, we have

$$\lim_{n \rightarrow \infty} m((x_n - x)([-\varepsilon, \varepsilon])) = 1$$

(ii) almost everywhere in the fuzzy state m , if for every $\varepsilon > 0$, we have

$$m\left(\bigvee_{k=1}^\infty \bigwedge_{n=k}^\infty ((x_n - x)([-\varepsilon, \varepsilon]))\right) = 1$$

(iii) everywhere, if

$$\bigwedge_{p=1}^\infty \bigvee_{k=1}^\infty \bigwedge_{n=k}^\infty \left((x_n - x)\left(\left[-\frac{1}{p}, \frac{1}{p}\right]\right) \right) = 1_{\mathbb{A}}$$

(iv) in a mean p , where $1 \leq p < \infty$, if

$$\lim_{n \rightarrow \infty} (m|\bar{x}_n - \bar{x}|^p) = \bar{0}$$

(v) everywhere on f , if

$$f \leq \bigwedge_{p=1}^\infty \bigvee_{k=1}^\infty \bigwedge_{n=k}^\infty \left((x_n - x)\left(\left[-\frac{1}{p}, \frac{1}{p}\right]\right) \right)$$

(vi) uniformly on $\bar{f} \in \mathcal{M}$, if for every $\varepsilon > 0$, there is an integer n_0 , such that

$$\forall n \geq n_0 : (x_n - x)([-\varepsilon, \varepsilon]) \geq \bar{f}$$

(vii) uniformly, if for every $\varepsilon > 0$, there is an integer n_0 , such that

$$\forall n \geq n_0 : (x_n - x)([-\varepsilon, \varepsilon]) = \bar{1}$$

(viii) almost uniformly in fuzzy state m , if for every $\epsilon > 0$ there is an element $f \in \mathcal{M}$, such that $m(f') \leq \epsilon$ and a sequence $\{x_n\}_{n=1}^\infty$ converges uniformly to x on f .

To prove the law of large numbers and the central limit theorem on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$, we also need the next theorem, which was proved in Dvurečenskij and Tírpaková [57].

Theorem 1. Let m be a fuzzy state of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$, x, x_1, x_2, \dots be fuzzy observables of $(\mathbb{A}, \mathcal{M})$, and u, u_1, u_2, \dots be functions with properties (1) and (2). Then,

- (A) The sequence of fuzzy observables $\{x_n\}_{n=1}^\infty$ converges to fuzzy observable x
 - (i) in fuzzy state m if and only if the sequence of functions $\{u_n\}_{n=1}^\infty$ converges to u in measure μ_φ ,
 - (ii) almost uniformly in fuzzy state m if and only if the sequence of functions $\{u_n\}_{n=1}^\infty$ converges almost uniformly to u in measure μ_φ ,
 - (iii) almost everywhere in fuzzy state m if and only if the sequence of functions $\{u_n\}_{n=1}^\infty$ converges almost everywhere to u in measure μ_φ ,
 - (iv) in mean p ($1 \leq p \leq \infty$) if and only if $\{u_n\}_{n=1}^\infty$ converges to u in mean p in measure μ_φ .
- (B) If the sequence of fuzzy observables $\{x_n\}_{n=1}^\infty$ converges to fuzzy observable x
 - (v) everywhere, then there is $\Lambda \in \mathcal{S}$ such that $\varphi(\Lambda) = \bar{1}_\mathbb{A}$ and the sequence $\{u_n\}_{n=1}^\infty$ converges to u everywhere on Λ ,
 - (vi) uniformly, then there is $\Lambda \in \mathcal{S}$ such that $\varphi(\Lambda) = \bar{1}_\mathbb{A}$ and the sequence $\{u_n\}_{n=1}^\infty$ converges to u uniformly on Λ ,
 - (vii) uniformly on $f \in \mathcal{M}$, then there is $\Lambda \in \mathcal{S}$ such that $\varphi(\Lambda) \geq \bar{f}$ and the sequence $\{u_n\}_{n=1}^\infty$ converges to u uniformly on Λ .

Conversely, if the sequence of functions $\{u_n\}_{n=1}^\infty$ defined by (1), (2) converges to u

- (viii) everywhere, then $\{x_n\}_{n=1}^\infty$ converges to fuzzy observable x everywhere on $f \in \mathcal{M}$, where $\bar{f} = \bar{1}_\mathbb{A}$,
- (ix) uniformly, then $\{x_n\}_{n=1}^\infty$ converges to fuzzy observable x uniformly on $f \in \mathcal{M}$, where $\bar{f} = \bar{1}_\mathbb{A}$
- (x) uniformly on $\Lambda, \Lambda \in \mathcal{S}$, then fuzzy observables $\{x_n\}_{n=1}^\infty$ converges to fuzzy observable x uniformly on $f \in \mathcal{M}$, where $\bar{f} = \varphi(\Lambda)$.

In the following, we will continue to introduce the notion of the independence of fuzzy observables $\{x_n\}_{n=1}^\infty$ in fuzzy state m . Now, we define the joint fuzzy observable of fuzzy observables.

Definition 6. Let $x_1, x_2, \dots, x_n, n \geq 2$, be a finite system of fuzzy observables on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$. A joint fuzzy observable of fuzzy observables x_1, x_2, \dots, x_n is a σ -homomorphism $T_n : \mathcal{B}(\mathbb{R}^n) \rightarrow \mathcal{M}$, such that

- (i) $T_n(A^c) = T_n(A)'$ for every $A \in \mathcal{B}(\mathbb{R}^n)$
- (ii) $T_n(\cup_{i=1}^n A_i) = \wedge_{i=1}^n T_n(A_i), A_i \in \mathcal{B}(\mathbb{R}^n), i = 1, 2, \dots, n$
- (iii) $T_n(\pi_i^{-1}(E)) = x_i(E)$ for every $i \in \{1, 2, \dots, n\}, E \in \mathcal{B}(\mathbb{R}^1)$

where $\pi_i : \mathbb{R}^n \rightarrow \mathbb{R}^1$ is the projection into the i -th coordinate.

In accordance with Riečan [61] and Riečan and Neubrunn [62], a sufficient condition for the existence of the joint fuzzy observable of fuzzy observables $x_1, x_2, \dots, x_n, n \geq 2$, meets the condition $x_i(\emptyset) = x_j(\emptyset)$ for every $i, j \in \{1, 2, \dots, n\}$.

Definition 7. Fuzzy observables x_1, x_2, \dots, x_n on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ are independent in fuzzy state m if for every $n \geq 2$ there exists joint fuzzy observable T_n and

$$m(T_n(E_1 \times E_2 \times \dots \times E_n)) = \prod_{i=1}^n m(x_i(E_i))$$

for any $E_i \in \mathcal{B}(\mathbb{R}^1), i = 1, 2, \dots, n$.

According to the assumption of independence of the sequence of fuzzy observables $\{x_n\}_{n=1}^\infty$, for every $n \geq 2$ there exists joint fuzzy observable T_n . To each fuzzy observable $x_i : \mathcal{B}(\mathbb{R}^1) \rightarrow \mathcal{M}, i = 1, 2, \dots, n$ exists the observable $\bar{x}_i = h \circ x_i : \mathcal{B}(\mathbb{R}^1) \rightarrow \mathcal{M} / \sim_m$ and a real function $u_i : \Omega \rightarrow \mathbb{R}^1$, such that $\bar{x}_i(E) = \varphi(u_i^{-1}(E))$. We define function $\Phi : \Omega \rightarrow \mathbb{R}^1$, such that $\Phi_n(\omega) = (u_1(\omega), u_2(\omega), \dots, u_n(\omega)), \omega \in \Omega$. If $\bar{T}_n = h \circ T_n : \mathcal{B}(\mathbb{R}^n) \rightarrow \mathcal{M} / \sim_m$, then $\bar{T}_n = \varphi \circ \Phi_n^{-1}$. The main idea of the proof can be illustrated by Figure 1.




Figure 1. The main idea of the proof of the central limit theorem.

4. Limit Theorems for Fuzzy Quantum Space

Theorem 2 (Central limit theorem). Let $\{x_n\}_{n=1}^\infty$ be a sequence of independent fuzzy observables, identically distributed in fuzzy state m , with mean value a and variance $\sigma^2 \in (0, \infty)$. Then, for any $s \in \mathbb{R}^1$, the following equality holds:

$$\lim_{n \rightarrow \infty} m \left(\frac{1}{\sigma \sqrt{n}} \sum_{i=1}^n (x_i - na)(-\infty, s) \right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^s e^{-\frac{t^2}{2}} dt$$

Proof. We define the real function $k_n : \mathbb{R}^n \rightarrow \mathbb{R}^1$ as follows

$$k_n(r_1, r_2, \dots, r_n) = \frac{1}{\sigma \sqrt{n}} \sum_{i=1}^n (r_i - na), r_i \in \mathbb{R}, i = 1, 2, \dots, n.$$

Calculate:

$$\begin{aligned} m \left(\frac{1}{\sigma \sqrt{n}} \sum_{i=1}^n (x_i - na)(-\infty, s) \right) &= m(T_n(k_n^{-1}((-\infty, s)))) = \mu(\bar{T}_n(k_n^{-1}((-\infty, s)))) = \\ &= \mu(\varphi \circ \Phi_n^{-1}(k_n^{-1}((-\infty, s)))) = \mu_\varphi(\Phi_n^{-1}(k_n^{-1}((-\infty, s)))) = \mu_\varphi(\{\omega : k_n(\Phi_n(\omega)) < s\}), \end{aligned}$$

where

$$k_n(\Phi_n(\omega)) = \frac{1}{\sigma\sqrt{n}} \sum_{i=1}^n (u_i - na).$$

Regarding Theorem 1.7.5 [63], the validity of the following argument is obvious:

$$\lim_{n \rightarrow \infty} m\left(\frac{1}{\sigma\sqrt{n}} \sum_{i=1}^n (x_i - na)(-\infty, s)\right) = \lim_{n \rightarrow \infty} \mu_\varphi(\{\omega : k_n(\Phi_n(\omega)) < s\}) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^s e^{-\frac{t^2}{2}} dt. \square$$

Theorem 3 (Weak law of large numbers). Let $\{x_n\}_{n=1}^\infty$ be a sequence of independent fuzzy observables, identically distributed in fuzzy state m , with the mean value a . Then,

$$\frac{x_1 + x_2 + \dots + x_n}{n} - a$$

converges to null fuzzy observable o in fuzzy state m .

Proof. We define real function $k_n : \mathbb{R}^n \rightarrow \mathbb{R}^1$ as follows

$$k_n(r_1, r_2, \dots, r_n) = \frac{1}{n} \sum_{i=1}^n r_i - a, r_i \in \mathbb{R}, i = 1, 2, \dots, n.$$

If T_n is the joint fuzzy observable of fuzzy observables x_1, x_2, \dots, x_n , then we define fuzzy observable $y_n = T_n \circ k_n^{-1}$. According to Definition 5 (i) a sequence of fuzzy observables $\{y_n\}_{n=1}^\infty$ converges to null fuzzy observable o in fuzzy state m , if for every $\varepsilon > 0$ it holds that

$$\sum_{i=1}^\infty m(y_n(-\varepsilon, \varepsilon)) = 1.$$

Calculate:

$$\begin{aligned} m(y_n((-\varepsilon, \varepsilon))) &= m(T_n(k_n^{-1}((-\varepsilon, \varepsilon)))) = \mu(\overline{T_n}(k_n^{-1}((-\varepsilon, \varepsilon)))) = \\ &= \mu(\varphi \circ \Phi_n^{-1}(k_n^{-1}((-\varepsilon, \varepsilon)))) = \mu_\varphi(\Phi_n^{-1}(k_n^{-1}((-\varepsilon, \varepsilon)))). \end{aligned}$$

If $k_n(\Phi_n(\omega)) = \frac{1}{n} \sum_{i=1}^n u_i(\omega) - a$, then according to Theorem 1 A) (i) the sequence of fuzzy observables $\{y_n\}_{n=1}^\infty$ converges to null fuzzy observable o in fuzzy state m if and only if the sequence of the functions $\{k_n(\Phi_n)\}_{n=1}^\infty$ converges to null in measure μ_φ . According to Theorem 1.10.3 [61], the validity of the arguments is obvious. \square

Theorem 4 (Strong law of large numbers). Let $\{x_n\}_{n=1}^\infty$ be a sequence of fuzzy observables independent in fuzzy state m , such that

$$\sum_{i=1}^\infty \frac{1}{i^2} D(x_i) < \infty.$$

Then,

$$\frac{1}{n} \sum_{i=1}^n (x_i - m(x_i))$$

converges to null fuzzy observable o almost everywhere in fuzzy state m .

Proof. We define real function $q_n : \mathbb{R}^n \rightarrow \mathbb{R}^1$ as follows:

$$q_n(r_1, r_2, \dots, r_n) = \frac{1}{n}(r_1 - E(r_1) + r_2 - E(r_2) + \dots + r_n - E(r_n)),$$

where $E(r_i)$ is the mean value r_i defined as $E(r_i) = \int_{\mathbb{R}^1} t d\mu_\varphi(t), r_i \in \mathbb{R}, i = 1, 2, \dots, n$. Then, $q_n(\Phi_n(\omega)) = \frac{1}{n}(\sum_{i=1}^n (u_i(\omega) - E(u_i(\omega))))$. If T_n is a joint fuzzy observable of fuzzy observables x_1, x_2, \dots, x_n , then we define fuzzy observable $z_n = T_n \circ q_n^{-1} = \frac{1}{n}(\sum_{i=1}^n (x_i - m(x_i)))$. According to Definition 5 (ii), a sequence of fuzzy observables $\{z_n\}_{n=1}^\infty$ converges to null fuzzy observable o almost everywhere in fuzzy state m if for any $\varepsilon > 0$ holds:

$$m(\bigvee_{k=1}^\infty \bigwedge_{n=k}^\infty (z_n([- \varepsilon, \varepsilon]))) = 1.$$

Then, according to Theorem 1 A) (iii), a sequence of fuzzy observables $\{z_n\}_{n=1}^\infty$ converges to null fuzzy observable o almost everywhere in fuzzy state m if and only if the sequence of functions $\{q_n(\Phi_n)\}_{n=1}^\infty$ converges to null almost everywhere in measure μ_φ . According to Theorem 1.10.5 [61], the validity of the arguments is obvious. \square

5. Extreme Value Theorems for Fuzzy Quantum Space

Let $\{x_n\}_{n=1}^\infty$ be a sequence of independent, identically-distributed fuzzy observables of fuzzy quantum space $(\mathbb{A}, \mathcal{M})$. For any $n \geq 1$ we define the real function $k_n : \mathbb{R}^n \rightarrow \mathbb{R}^1$ as follows:

$$k_n(r_1, r_2, \dots, r_n) = \max\{r_1, r_2, \dots, r_n\}.$$

Let T_n be the joint fuzzy observable of fuzzy observables x_1, x_2, \dots, x_n . We define the maximum fuzzy observables of x_1, x_2, \dots, x_n as

$$M_1 = x_1, M_n = \max\{x_1, x_2, \dots, x_n\} = T_n \circ k_n^{-1}, n \geq 2,$$

where M_n is the fuzzy observable.

Theorem 5. Let $(\xi_n)_{n=1}^\infty$ be a sequence of independent random variables with the same distribution function $F : \mathbb{R} \rightarrow [0, 1]$ in fuzzy state m . Put $\eta_n = \max(\xi_1, \dots, \xi_n), n = 1, 2, \dots$. Let there exist $a_n > 0, b_n \in \mathbb{R}$ such that

$$\lim_{n \rightarrow \infty} m\left(\left\{\omega; \frac{\eta_n(\omega) - b_n}{a_n} < x\right\}\right) = \lim_{n \rightarrow \infty} F^n(a_n x + b_n) = H(x),$$

where $H : \mathbb{R} \rightarrow [0, 1]$ is a continuous distribution function, increasing on an interval. Then, H has one of three distributions with parameters $\mu, \sigma, \alpha > 0$ (Figure 2):

1. Gumbel $H_{\mu, \sigma}(x) = \exp\left(-e^{-\left(\frac{x-\mu}{\sigma}\right)}\right), x \in \mathbb{R}$
2. Fréchet $H_{\mu, \sigma, \alpha}(x) = \begin{cases} 0, & \text{for } x \leq \mu \\ \exp\left(-\left(\frac{x-\mu}{\sigma}\right)^{-\alpha}\right), & \text{for } x > \mu, \alpha > 0 \end{cases}$
3. Weibull $H_{\mu, \sigma, \alpha}(x) = \begin{cases} \exp\left(-\left(-\frac{x-\mu}{\sigma}\right)^\alpha\right), & \text{for } x \leq \mu, \alpha > 0 \\ 1, & \text{for } x > \mu \end{cases}$




Figure 2. Densities of the standard extreme value distribution .

Proof of Theorem 5 can be found in [63].

Theorem 6 (Fisher–Tippett–Gnedenko theorem). Let $\{x_n\}_{n=1}^\infty$ be a sequence of fuzzy independent observables identically distributed in fuzzy state m . Let there exist norming constants $a_n > 0, b_n \in \mathbb{R}$ and some non-degenerate distribution function H such that

$$\lim_{n \rightarrow \infty} m\left(\frac{1}{a_n}(M_n - b_n)(-\infty, t)\right) = H(t) \text{ for any } t \in \mathbb{R}.$$

Then, H belongs to the type of one of the following three types of standard extreme value distributions: Gumbel, Fréchet, or Weibull.

Proof. We define the real function $q_n : \mathbb{R}^n \rightarrow \mathbb{R}^1$ as follows:

$$q_n(r_1, r_2, \dots, r_n) = \frac{1}{a_n}(\max\{r_1, r_2, \dots, r_n\} - b_n), \quad r_i \in \mathbb{R}^1, \quad i = 1, 2, \dots, n.$$

Then $q_n(\Phi_n(\omega)) = \frac{1}{a_n}(\max\{u_1(\omega), u_2(\omega), \dots, u_n(\omega)\} - b_n)$ and

$$\begin{aligned} m\left(\frac{1}{a_n}(M_n - b_n)(-\infty, t)\right) &= m(T_n(q_n^{-1}((-\infty, t)))) = \mu(\overline{T}_n(q_n^{-1}((-\infty, t)))) = \\ &= \mu(\varphi \circ \Phi_n^{-1}(q_n^{-1}((-\infty, t)))) = \mu_\varphi(\Phi_n^{-1}(q_n^{-1}((-\infty, t)))) = \mu_\varphi(\{\omega : q_n(\Phi_n(\omega)) < t\}). \end{aligned}$$

We have

$$H(t) = \lim_{n \rightarrow \infty} m\left(\frac{1}{a_n}(M_n - b_n)(-\infty, t)\right) = \lim_{n \rightarrow \infty} \mu_\varphi(\{\omega : q_n(\Phi_n(\omega)) < t\}).$$

Then, according to Theorem 5, the validity of the arguments is obvious. \square

Now, we define the distribution function and excess distribution function on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$.

Definition 8. Let $m : \mathcal{M} \rightarrow [0, 1]$ be a fuzzy state and $x : \mathcal{B}(\mathbb{R}^1) \rightarrow \mathcal{M}$ be a fuzzy observable on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$. For any $t \in \mathbb{R}^1$ we define function $F_x : \mathbb{R}^1 \rightarrow [0, 1]$ as

$$F_x(t) = m(x((-\infty, t))).$$

Function F_x is called the distribution function of an observable x on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$.

Proposition 1. If the function F_x is the distribution function of an observable x on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$, then it satisfies the following conditions:

- (i) F_x is non-decreasing
- (ii) F_x is left continuous
- (iii) $\lim_{n \rightarrow \infty} F_x = 1$
- (iv) $\lim_{n \rightarrow -\infty} F_x = 0$

Proof.

- (i) Let $s < t, s, t \in \mathbb{R}^1$, then $x((-\infty, s)) \leq x((-\infty, t))$, it follows that

$$F_x(s) = m(x((-\infty, s))) \leq m(x((-\infty, t))) = F_x(t).$$

We proved that the function F_x is non-decreasing.

- (ii) Let $t_n \nearrow t, t_n, t \in \mathbb{R}^1, n = 1, 2, \dots$, then $x((-\infty, t_n)) \nearrow x((-\infty, t))$, it follows that

$$F_x(t_n) = m(x((-\infty, t_n))) \nearrow m(x((-\infty, t))) = F_x(t).$$

We proved that the function F_x is left continuous.

- (iii) Let $t_n \nearrow \infty, t_n \in \mathbb{R}^1, n = 1, 2, \dots$, then $x((-\infty, t_n)) \nearrow x((-\infty, \infty)) = 1_{\mathbb{A}}$, it follows that

$$F_x(t_n) = m(x((-\infty, t_n))) \nearrow m(x((-\infty, \infty))) = m(1_{\mathbb{A}}) = 1.$$

We proved that $\lim_{n \rightarrow \infty} F_x = 1$.

- (iv) Let $t_n \searrow -\infty, t_n \in \mathbb{R}^1, n = 1, 2, \dots$, then $x((-\infty, t_n)) \searrow x((-\infty, -\infty)) = 0_{\mathbb{A}}$, it follows that

$$F_x(t_n) = m(x((-\infty, t_n))) \searrow m(x((-\infty, -\infty))) = m(0_{\mathbb{A}}) = 0.$$

We proved that $\lim_{n \rightarrow -\infty} F_x = 0$. \square

Now we define the excess distribution function \tilde{F}_w on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$.

Definition 9. For $w > 0$ we define excess distribution function \tilde{F}_w on fuzzy quantum space $(\mathbb{A}, \mathcal{M})$ as

$$\tilde{F}_w(t) = \frac{\tilde{F}(t+w) - \tilde{F}(w)}{1 - \tilde{F}(w)}$$

for every $0 < t < \omega(\tilde{F}) = \sup\{t; \tilde{F}(t) < 1\}$. Value $\omega(\tilde{F})$ is called the right endpoint of distribution function \tilde{F} .

Theorem 7 (Balkema, de Haan–Pickands). For a sufficiently large w , the excess distribution \tilde{F}_w converges to the generalized pareto distribution. Parameter $\beta = \beta(w)$ is dependent on threshold w , and for every $\alpha > 0$

$$\lim_{w \rightarrow \omega(F_x)} \sup_{0 \leq t \leq \omega(F_x) - w} |\tilde{F}_w(t) - G_{\alpha, \beta(w)}(t)| = 0.$$

Proof. Let $x_i : \mathcal{B}(\mathbb{R}^1) \rightarrow M, i = 1, 2, \dots, n$ be fuzzy observables. Then, there exist observables $\bar{x}_i = h \circ x_i : \mathcal{B}(\mathbb{R}^1) \rightarrow \mathcal{M}/I_m$ and real functions $u_i : \Omega \rightarrow \mathbb{R}^1$, such that $\bar{x}_i(E) = \varphi(u_i^{-1}(E))$. Then,

$$\begin{aligned} \tilde{F}(t) &= m(x((-\infty, t))) = \mu(\bar{x}_i((-\infty, t))) = \mu(\varphi \circ u_n^{-1}((-\infty, t))) = \\ &= \mu_\varphi(u_n^{-1}((-\infty, t))) = \mu_\varphi(\{\omega : u_n(\omega) < t\}) = F(t), t \in \mathbb{R} \end{aligned}$$

is the distribution function of real random variable u . It is obvious that

$$\tilde{F}_w(t) = \frac{\tilde{F}(t+w) - \tilde{F}(w)}{1 - \tilde{F}(w)} = \frac{F(t+w) - F(w)}{1 - F(w)} = F_w(t).$$

□

6. Conclusions

The seminal theoretical results in probability theory are limit theorems. When using random samples to estimate distributional parameters, we would like to know that as the sample size gets larger, the estimates are probably close to the parameters that they are estimating. In statistical inference, the central limit theorem is the dominant and most useful theorem. It allows us to make the assumption that, for a population, a normal distribution will occur regardless of what the initial distribution looks like for a sufficiently large sample size. When the distribution shape is not known or the population is not normally distributed, the theorem is used to make assumptions. The law of large numbers is an invaluable tool that is expected to state definite things about the real-world results of unexpected events. The law of large numbers is the postulate of statistics and probability theory that states that the greater the number of samples are used from an event, the closer the monitoring results will be to the average population. Thus, the law of large numbers describes the stability of big random variables. Both the strong and weak laws refer to the convergence of the sample mean to the population mean as the sample size gets bigger.

This paper generalizes the central limit theorem, the law of large numbers, and extreme value theorems of classical probability theory to fuzzy quantum spaces. Extreme value theory models rare events outside the range of allowable observations with high impact. This method has become a widely-used tool for risk assessment in recent years. It is used in the areas of insurance, banking, operational risk, market risk, and credit risk [52]. By applying these limit theorems to the Atanassov set, it gives us the space to work with incomplete data, which we can use in the area of finance. The basic advantage of fuzzy logic is the ability to mathematically express information expressed verbally. Thanks to this, fuzzy logic proves to be a very good tool for working with behavioral data. Behavioral finance takes into account the human factor when making financial decisions. For this reason, behavioral finance often uses linguistic data, and therefore it is appropriate to use methods based on fuzzy logic to describe them. Behavioral finance is a financial field examining the effect of social, cognitive, and emotional factors on the economic decisions of individuals and institutions as well as the consequences of these decisions on market prices [64].

Author Contributions: All authors contributed equally and significantly in writing this article. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: Our thanks belong to Beloslav Riečan, who contributed the results used in our work. Rest in peace, our beloved friend, co-worker, and teacher.

Conflicts of Interest: The authors declare that they have no conflicts of interests.

References

- Kolmogorov, A.N. *Basic Probability Concepts*; Nauka: Moscow, Russia, 1974; p. 119.
- Čunderlíková, K.; Bartková, R. Basic Theorems from Extreme Value Theory for MV-algebras. In *Issues in Intuitionistic Fuzzy Sets and Generalized Nets*; Polish Academy of Sciences: Warsaw, Poland, 2020; Volume 14, ISBN 9788361551232.
- Birkhoff, G.; von Neumann, J. The logic of quantum mechanics. *Ann. Math.* **1936**, *37*, 823–834. [[CrossRef](#)]
- Heisenberg, W. Über quantentheoretische Umdeutung kinematischer und mechanischer Beziehungen. *Z. Phys.* **1925**, *33*, 879–893. [[CrossRef](#)]
- Schrödinger, E. Quantisierung als Eigenwertproblem. *Ann. Phys.* **1926**, *384*, 361–376. [[CrossRef](#)]
- Zadeh, L.A. Probability measures of fuzzy events. *J. Math. Anal. Appl.* **1968**, *23*, 421–427. [[CrossRef](#)]
- Varadarajan, V.S. *Geometry of Quantum Theory*, 2nd ed.; Springer: New York, NY, USA, 2007; ISBN 9780387961248.
- Piasecki, K. Probability of fuzzy events defined as denumerable additivity measure. *Fuzzy Sets Syst.* **1985**, *17*, 271–284. [[CrossRef](#)]
- Riečan, B. A new approach to some notions of statistical quantum mechanics. *Busefal* **1988**, *35*, 4–6.
- Pykacz, J. Quantum logics and soft fuzzy probability spaces. *Busefal* **1987**, *32*, 150–157.
- Dvurečenskij, A.; Chovanec, F. Fuzzy quantum spaces and compatibility. *Int. J. Theor. Phys.* **1988**, *27*, 1069–1082. [[CrossRef](#)]
- Dvurečenskij, A. On the existence of probability measures on fuzzy measurable spaces. *Fuzzy Sets Syst.* **1991**, *43*, 173–181. [[CrossRef](#)]
- Navara, M. Existence of states on quantum structures. *Inf. Sci.* **2009**, *179*, 508–514. [[CrossRef](#)]
- Navara, M.; Pták, P. P-measures on soft fuzzy σ -algebras. *Fuzzy Sets Syst.* **1993**, *56*, 123–126. [[CrossRef](#)]
- Navara, M.; Pták, P. On the state space of soft fuzzy algebras. *Busefal* **1991**, *48*, 55–63.
- Dvurečenskij, A.; Riečan, B. On joint distribution of observables for F-quantum spaces. *Fuzzy Sets Syst.* **1991**, *20*, 65–73. [[CrossRef](#)]
- Dvurečenskij, A.; Riečan, B. Fuzzy quantum models. *Int. J. Gen. Syst.* **1991**, *20*, 39–54. [[CrossRef](#)]
- Dvurečenskij, A.; Kôpka, F.; Riečan, B. On a representation theorem of observables in ordered spaces. *Busefal* **1993**, *56*, 15–19.
- Riečan, B. On mean value in F-quantum spaces. *Appl. Math.* **1990**, *35*, 209–214. [[CrossRef](#)]
- Riečan, B. Indefinite integral in fuzzy quantum spaces. *Busefal* **1989**, *38*, 5–7.
- Mesiar, R. Fuzzy probability measures and the Bayes formula. *Busefal* **1993**, *56*.
- Mesiar, R. Symmetric fuzzy probability measures and the Bayes formula. *Busefal* **1988**, *34*, 130–137.
- Mesiar, R. Fuzzy Bayes formula and the ill defined elements. In *Proceedings of the First Winter School on Measure Theory, Bratislava, Slovakia, 10–15 January 1988*; pp. 82–87.
- Piasecki, K. Note on “On the Bayes formula for fuzzy probability measure”. *Fuzzy Sets Syst.* **1987**, *24*, 121–122. [[CrossRef](#)]
- Piasecki, K. On the Bayes formula for fuzzy probability measure. *Fuzzy Sets Syst.* **1986**, *18*, 183–185. [[CrossRef](#)]
- Piasecki, L.; Svitalski, Z. A remark on the definition of fuzzy P-measure and the Bayes formula. *Fuzzy Sets Syst.* **1988**, *26*, 112–124. [[CrossRef](#)]
- Markechová, D. The entropy on F-quantum spaces. *Math. Slovaca* **1993**, *40*, 177–190.
- Markechová, D. Isomorphism and conjugation of fuzzy dynamical systems. *Busefal* **1989**, *38*, 94–101.
- Markechová, D. The entropy of fuzzy dynamical systems. *Busefal* **1989**, *38*, 38–41.
- Tirpakova, A.; Markechova, D. The fuzzy analogies of some ergodic theorems. *Adv. Differ. Equ.* **2015**, *2015*, 1–10. [[CrossRef](#)]
- Markechová, D.; Tirpáková, A. Birkhoff’s individual ergodic theorem and maximal ergodic theorem for fuzzy dynamical systems. *Adv. Differ. Equ.* **2016**, 1–8. [[CrossRef](#)]
- Harman, B.; Riečan, B. On the individual ergodic theorem in F-quantum spaces. *Zesz. Nauk. Akad. Ekon. Pozn. Ser. 1* **1992**, *187*, 25–30.
- Dvurečenskij, A.; Pulmanová, S. On joint distribution of observables. *Math. Slovaca* **1982**, *32*, 155–166.
- Jajte, R. Strong laws of large numbers for several contractions in a von Neumann algebra. *Lect. Notes Math.* **1989**, *1391*, 125–139.
- Ochs, W. On strong law of large numbers in quantum probability theory. *J. Philos. Log.* **1977**, *6*, 473–480. [[CrossRef](#)]
- Ochs, W. Concepts of convergence for a quantum law of large numbers. *Rep. Math. Phys.* **1980**, *17*, 127–143. [[CrossRef](#)]
- Cushen, C.D.; Hudson, R.L. A quantum mechanical central limit theorem. *J. Appl. Probab.* **1971**, *8*, 454–469. [[CrossRef](#)]
- Gudder, S. Joint distributions of observables. *J. Math. Mech.* **1968**, *18*, 325–335. [[CrossRef](#)]
- Révész, P. *The Laws of Large Numbers*; Academic Press: Cambridge, MA, USA, 1968; ISBN 0125871503.
- Dvurečenskij, A. The Radon-Nikodým theorem for fuzzy probability spaces. *Fuzzy Sets Syst.* **1992**, *45*, 69–78. [[CrossRef](#)]
- Riečan, B. On the convergence of observables in fuzzy quantum logics. *Tatra Mountains Math. Publ.* **1995**, *6*, 149–156.
- Riečan, B. On some mathematical models of quantum mechanical systems. *Busefal* **1993**, *56*, 63–66.
- Chovanec, F.; Kôpka, F. Fuzzy equality and convergences for F-observables in F-quantum spaces. *Appl. Math.* **1991**, *36*, 32–45. [[CrossRef](#)]
- Kôpka, F.; Chovanec, F. Martingale convergence theorem in F-quantum spaces. *Busefal* **1993**, *56*, 20–28.
- Fisher, R.A.; Tippett, L.H.C. Limiting Forms of the Frequency Distribution of the Largest or Smallest Members of a Sample. *Proc. Camb. Philos. Soc.* **1928**, *24*, 180–190. [[CrossRef](#)]

46. Gnedenko, B. Sur La Distribution Limite Du Terme Maximum D'Une Serie Aleatoire. *Ann. Math.* **1943**, *44*, 423–453. [[CrossRef](#)]
47. Kotz, S.; Nadarajah, S. *Extreme Value Distributions: Theory and Applications*; Imperial College Press: London, UK, 2000. [[CrossRef](#)]
48. David, H.A.; Edwards, A.W.F. *Annotated Readings in the History of Statistics*; Springer: New York, NY, USA, 2001; ISBN 9781441931740.
49. Feller, W. *An Introduction to Probability Theory and Its Applications II*, 2nd ed.; John Wiley & Sons: Hoboken, NJ, USA, 1971.
50. Garrido, M.C.; Lezaud, P. Extreme Value Analysis: An Introduction. *J. Soc. Française Stat.* **2013**, *154*, 66–97.
51. Coles, S. *An Introduction to Statistical Modeling of Extreme Values*; Springer: New York, NY, USA, 2001; ISBN 9781447136750.
52. Embrechts, P.; Kluppelberg, C.; Mikosch, T. *Modelling Extremal Events for Insurance and Finance*; Springer: New York, NY, USA, 1997; Volume 180, pp. 152–180. ISSN 01724568.
53. Gudder, S.; Mullikin, H.C. Measure theoretic convergences of observables and operators. *J. Math. Phys.* **1973**, *14*, 234–242. [[CrossRef](#)]
54. Haan, L.; Ferreira, A. *Extreme Value Theory: An Introduction*; Springer: New York, NY, USA, 2006; ISBN 9780387344713.
55. Markechová, D. *F-Dynamical Systems and Their Entropy*; UKF: Nitra, Slovakia, 2011; p. 80. ISBN 9788080948849.
56. Markechová, D. *Fuzzy Probability Spaces, Fuzzy Dynamical Systems, and Entropy*; Lambert Academic Publishing: Riga, Latvia, 2013; p. 107. ISBN 9783659463341.
57. Dvurečenskij, A.; Tírpáková, A. Sum of observables in fuzzy quantum spaces. *Appl. Math.* **1992**, *37*, 40–50. [[CrossRef](#)]
58. Riečan, B. Probability Theory on IF Events. In *Algebraic and Proof-Theoretic Aspects of Non-Classical Logics*; Lecture Notes in Computer Science; Aguzzoli, S., Ciabottoni, A., Gerla, B., Manara, C., Marra, V., Eds.; Springer: Berlin/Heidelberg, Germany, 2007; Volume 4460, ISBN 9783540759386. [[CrossRef](#)]
59. Sikorsk, R. Boolean algebras. In *Ergebnisse der Mathematik und Ihre Grenzgebiete*, 3rd ed.; Springer: Berlin/Heidelberg, Germany, 1964; p. 25.
60. Dvurečenskij, A. Loomis-Sikorski theorem for σ -complete MV-algebras and ℓ -groups. *J. Aust. Math. Soc.* **2000**, *68*, 261–277. [[CrossRef](#)]
61. Riečan, B. Probability theory and the operations with IF-sets. In Proceedings of the World Congress on Computational Intelligence, Hong Kong, China, 1–6 June 2008; pp. 1250–1252. [[CrossRef](#)]
62. Riečan, B.; Neubrunn, T. *Measure Theory*; Veda: Bratislava, Slovakia, 1992; ISBN 9788022403689.
63. Bartková, R.; Riečan, B.; Tírpáková, A. *Probability Theory for Fuzzy Quantum Space with Statistical Applications*; Bentham Science Publishers: Soest, The Netherlands, 2017; p. 191. ISBN 9781681085395.
64. Ritter, J.R. Behavioral Finance. *Pac. Basin Financ. J.* **2003**, *11*, 429–437. [[CrossRef](#)]